



Dear 6<sup>th</sup> grade parents,

### **Vaccine requirements for 7<sup>th</sup> graders**

The state of Ohio requires all students entering the 7<sup>th</sup> grade to have the meningococcal vaccine (also known as MCV4, Menevo or Menactra) in addition to one dose of Tdap (Tetanus, diphtheria and pertussis).

#### **Meningococcal meningitis**

Meningococcal meningitis is a rare but serious disease that develops rapidly and can claim a life in as little as one day. Of those who survive, approximately one in five are left with serious medical problems like amputation, deafness, and brain damage. Teens are at increased risk of meningococcal meningitis. This increased risk may be due to activities like sharing utensils and kissing.

#### **Tetanus, diphtheria, and pertussis (Tdap)**

Tetanus causes painful tightening of the muscles, usually all over your body; diphtheria causes a thick covering in the back of the throat and can also lead to breathing problems, paralysis, heart failure, and even death; pertussis causes coughing spells and can lead to pneumonia, seizures, brain damage, and death, particularly in infants.

Vaccination is the best way to protect someone from getting the disease.

You may obtain these vaccines from your child's health care provider or other community resource. Vaccines are also available at your local health department. See school website for Summit County Health Department open hours.

Students who do not provide documentation of these immunizations to the school this fall are subject to exclusion. If there are medical/religious or philosophical reasons your child cannot receive immunizations, please contact the school clinic for an Immunization Exemption Form.

Thank you,

Tina Norris, Principal  
Coventry Middle School

Victoria Dorsey, District RN Supervisor for Coventry Local Schools  
Akron Children's Hospital, School Health Services

*See school website for related immunization information from the Centers for Disease Control (CDC)*

Talk to your child's doctor or nurse about the vaccines recommended for their age.

	Flu <i>Influenza</i>	Tdap Tetanus, diphtheria, pertussis	HPV Human papillomavirus	Meningococcal		Pneumococcal	Hepatitis B	Hepatitis A	Inactivated Polio	MMR Measles, mumps, rubella	Chickenpox <i>Varicella</i>
				MenACWY	MenB						
7-8 Years	Black	Light Gray		Light Gray		Light Gray	Light Gray	Light Gray	Light Gray	Light Gray	Light Gray
9-10 Years	Black	Light Gray	Light Gray	Light Gray		Light Gray	Light Gray	Light Gray	Light Gray	Light Gray	Light Gray
11-12 Years	Black	Black	Black	Black		Light Gray	Light Gray	Light Gray	Light Gray	Light Gray	Light Gray
13-15 Years	Black	Light Gray	Light Gray	Light Gray		Light Gray	Light Gray	Light Gray	Light Gray	Light Gray	Light Gray
16-18 Years	Black	Light Gray	Light Gray	Light Gray	Light Gray	Light Gray	Light Gray	Light Gray	Light Gray	Light Gray	Light Gray

**More information:**

Preteens and teens should get a flu vaccine every year.

Preteens and teens should get one shot of Tdap at age 11 or 12 years.

All 11-12 year olds should get a 2-shot series of HPV vaccine at least 6 months apart. A 3-shot series is needed for those with weakened immune systems and those age 15 or older.

All 11-12 year olds should get a single shot of a meningococcal conjugate (MenACWY) vaccine. A booster shot is recommended at age 16.

Teens, 16-18 years old, **may** be vaccinated with a serogroup B meningococcal (MenB) vaccine.

 These shaded boxes indicate when the vaccine is recommended for all children unless your doctor tells you that your child cannot safely receive the vaccine.

 These shaded boxes indicate the vaccine should be given if a child is catching-up on missed vaccines.

 These shaded boxes indicate the vaccine is recommended for children with certain health or lifestyle conditions that put them at an increased risk for serious diseases. See vaccine-specific recommendations at [www.cdc.gov/vaccines/pubs/ACIP-list.htm](http://www.cdc.gov/vaccines/pubs/ACIP-list.htm).

 This shaded box indicates children not at increased risk may get the vaccine if they wish after speaking to a provider.



U.S. Department of Health and Human Services  
Centers for Disease Control and Prevention

American Academy of Pediatrics



DEDICATED TO THE HEALTH OF ALL CHILDREN™



AMERICAN ACADEMY OF FAMILY PHYSICIANS  
STRONG MEDICINE FOR AMERICA

# Vaccine-Preventable Diseases and the Vaccines that Prevent Them

## **Diphtheria** (Can be prevented by Tdap vaccination)

Diphtheria is a very contagious bacterial disease that affects the respiratory system, including the lungs. Diphtheria bacteria can be spread from person to person by direct contact with droplets from an infected person's cough or sneeze. When people are infected, the bacteria can produce a toxin (poison) in the body that can cause a thick coating in the back of the nose or throat that makes it hard to breathe or swallow. Effects from this toxin can also lead to swelling of the heart muscle and, in some cases, heart failure. In serious cases, the illness can cause coma, paralysis, or even death.

## **Hepatitis A** (Can be prevented by HepA vaccination)

Hepatitis A is an infection in the liver caused by hepatitis A virus. The virus is spread primarily person-to-person through the fecal-oral route. In other words, the virus is taken in by mouth from contact with objects, food, or drinks contaminated by the feces (stool) of an infected person. Symptoms can include fever, tiredness, poor appetite, vomiting, stomach pain, and sometimes jaundice (when skin and eyes turn yellow). An infected person may have no symptoms, may have mild illness for a week or two, may have severe illness for several months, or may rarely develop liver failure and die from the infection. In the U.S., about 100 people a year die from hepatitis A.

## **Hepatitis B** (Can be prevented by HepB vaccination)

Hepatitis B causes a flu-like illness with loss of appetite, nausea, vomiting, rashes, joint pain, and jaundice. Symptoms of acute hepatitis B include fever, fatigue, loss of appetite, nausea, vomiting, pain in joints and stomach, dark urine, grey-colored stools, and jaundice (when skin and eyes turn yellow).

## **Human Papillomavirus** (Can be prevented by HPV vaccination)

Human papillomavirus is a common virus. HPV is most common in people in their teens and early 20s. About 14 million people, including teens, become infected with HPV each year. HPV infection can cause cervical, vaginal, and vulvar cancers in women and penile cancer in men. HPV can also cause anal cancer, oropharyngeal cancer (back of the throat), and genital warts in both men and women.

## **Influenza** (Can be prevented by annual flu vaccination)

Influenza is a highly contagious viral infection of the nose, throat, and lungs. The virus spreads easily through droplets when an infected person coughs or sneezes and can cause mild to severe illness. Typical symptoms include a sudden high fever, chills, a dry cough, headache, runny nose, sore throat, and muscle and joint pain. Extreme fatigue can last from several days to weeks. Influenza may lead to hospitalization or even death, even among previously healthy children.

## **Measles** (Can be prevented by MMR vaccination)

Measles is one of the most contagious viral diseases. Measles virus is spread by direct contact with the airborne respiratory droplets of an infected person. Measles is so contagious that just being in the same room after a person who has measles has already left can result in infection. Symptoms usually include a rash, fever, cough, and red, watery eyes. Fever can persist, rash can last for up to a week, and coughing can last about 10 days. Measles can also cause pneumonia, seizures, brain damage, or death.

## **Meningococcal Disease** (Can be prevented by meningococcal vaccination)

Meningococcal disease has two common outcomes: meningitis (infection of the lining of the brain and spinal cord) and bloodstream infections. The bacteria that cause meningococcal disease spread through the exchange of nose and throat droplets, such as when coughing, sneezing, or kissing. Symptoms include sudden onset of fever, headache, and stiff neck. With bloodstream infection, symptoms also include a dark purple rash. About one of every ten people who gets the disease dies from it. Survivors of meningococcal disease may lose their arms or legs, become deaf, have problems with their nervous systems, become developmentally disabled, or suffer seizures or strokes.

## **Mumps** (Can be prevented by MMR vaccination)

Mumps is an infectious disease caused by the mumps virus, which is spread in the air by a cough or sneeze from an infected person. A child can also get infected with mumps by coming in contact with a contaminated object, like a toy. The mumps virus causes swollen salivary glands under the ears or jaw, fever, muscle aches, tiredness, abdominal pain, and loss of appetite. Severe complications for children who get mumps are uncommon, but can include meningitis (infection of the covering of the brain and spinal cord), encephalitis (inflammation of the brain), permanent hearing loss, or swelling of the testes, which rarely results in decreased fertility.

## **Pertussis** (Whooping Cough) (Can be prevented by Tdap vaccination)

Pertussis spreads very easily through coughing and sneezing. It can cause a bad cough that makes someone gasp for air after coughing fits. This cough can last for many weeks, which can make preteens and teens miss school and other activities. Pertussis can be deadly for babies who are too young to receive the vaccine. Often babies get whooping cough from their older brothers or sisters, like preteens or teens, or other people in the family. Babies with pertussis can get pneumonia, have seizures, become brain damaged, or even die. About half of children under 1 year of age who get pertussis must be hospitalized.

## **Pneumococcal Disease** (Can be prevented by pneumococcal vaccination)

Pneumonia is an infection of the lungs that can be caused by the bacteria called pneumococcus. These bacteria can cause other types of infections too, such as ear infections, sinus infections, meningitis (infection of the lining of the brain and spinal cord), and bloodstream infections. Sinus and ear infections are usually mild and are much more common than the more serious forms of pneumococcal disease. However, in some cases pneumococcal disease can be fatal or result in long-term problems, like brain damage and hearing loss. The bacteria that cause pneumococcal disease spread when people cough or sneeze. Many people have the bacteria in their nose or throat at one time or another without being ill—this is known as being a carrier.

## **Polio** (Can be prevented by IPV vaccination)

Polio is caused by a virus that lives in an infected person's throat and intestines. It spreads through contact with the stool of an infected person and through droplets from a sneeze or cough. Symptoms typically include sore throat, fever, tiredness, nausea, headache, or stomach pain. In about 1% of cases, polio can cause paralysis. Among those who are paralyzed, about 2 to 10 children out of 100 die because the virus affects the muscles that help them breathe.

## **Rubella** (German Measles) (Can be prevented by MMR vaccination)

Rubella is caused by a virus that is spread through coughing and sneezing. In children rubella usually causes a mild illness with fever, swollen glands, and a rash that lasts about 3 days. Rubella rarely causes serious illness or complications in children, but can be very serious to a baby in the womb. If a pregnant woman is infected, the result to the baby can be devastating, including miscarriage, serious heart defects, mental retardation and loss of hearing and eye sight.

## **Tetanus** (Lockjaw) (Can be prevented by Tdap vaccination)

Tetanus mainly affects the neck and belly. When people are infected, the bacteria produce a toxin (poison) that causes muscles to become tight, which is very painful. This can lead to "locking" of the jaw so a person cannot open his or her mouth, swallow, or breathe. The bacteria that cause tetanus are found in soil, dust, and manure. The bacteria enter the body through a puncture, cut, or sore on the skin. Complete recovery from tetanus can take months. One to two out of 10 people who get tetanus die from the disease.

## **Varicella** (Chickenpox) (Can be prevented by varicella vaccination)

Chickenpox is caused by the varicella zoster virus. Chickenpox is very contagious and spreads very easily from infected people. The virus can spread from either a cough, sneeze. It can also spread from the blisters on the skin, either by touching them or by breathing in these viral particles. Typical symptoms of chickenpox include an itchy rash with blisters, tiredness, headache and fever. Chickenpox is usually mild, but it can lead to severe skin infections, pneumonia, encephalitis (brain swelling), or even death.

If you have any questions about your child's vaccines, talk to your healthcare provider.



# Meningococcal Vaccines for Preteens and Teens

Last updated NOVEMBER 2015

## Why does my child need to be vaccinated?

Meningococcal vaccines help protect against the bacteria that cause meningococcal disease. These infections don't happen very often, but can be very dangerous when they do. Meningococcal disease refers to any illness that is caused by *Neisseria meningitidis* bacteria. The two most severe and common illnesses caused by these bacteria include infections of the fluid and lining around the brain and spinal cord (meningitis) and bloodstream infections (bacteremia or septicemia). Even if they get treatment, about 10 to 15 out of 100 people with meningococcal disease will die from it.

Meningococcal disease can spread from person to person. The bacteria that cause this infection can spread when people have close or lengthy contact with someone's saliva, like through kissing or coughing, especially if they are living in the same place. Teens and young adults are at increased risk for meningococcal disease.

Meningococcal disease can become very serious, very quickly. The meningococcal vaccine is the best way to protect teens from getting meningococcal disease.

## When should my child be vaccinated?

All 11 to 12 year olds should be vaccinated with a single dose of a quadrivalent meningococcal conjugate vaccine. Older teens need a second shot when they are 16 years old so they stay protected when their risk is the highest.

Teens who got meningococcal vaccine for the first time when were 13, 14, or 15 years old should still get the booster shot when they are 16 years old. If your older teen didn't get the meningococcal shot at all, you should talk to their doctor about getting it as soon as possible.

Teens and young adults (16 through 23 year olds) may also be vaccinated with a serogroup B meningococcal vaccine (2 or 3 doses depending on brand), preferably at 16 through 18 years old. Talk with your teen's doctor or nurse about meningococcal vaccination to help protect your child's health.

## What else should I know about the vaccination?

Like many vaccines, meningococcal shots may cause mild side effects, like redness and soreness where the shot was given (usually in the arm). Note that both meningococcal vaccines can be given during the same visit, but in different arms. Some preteens and teens might faint after getting a meningococcal vaccine or any shot. To help avoid fainting, preteens and teens should sit or lie down when they get a shot and then for about 15 minutes after getting the shot.

## How can I get help paying for these vaccines?

The Vaccines for Children (VFC) program provides vaccines for children ages 18 years and younger, who are uninsured, Medicaid-eligible, American Indian or Alaska Native. You can find out more about the VFC program by going online to [www.cdc.gov](http://www.cdc.gov) and typing VFC in the search box.

## Where can I learn more?

Talk to your child's doctor or nurse to learn more about meningococcal vaccines and the other vaccines that your child may need. You can also find out more about these vaccines on CDC's Vaccines for Preteens and Teens website at [www.cdc.gov/vaccines/teens](http://www.cdc.gov/vaccines/teens).

DISTRIBUTED BY:



**U.S. Department of  
Health and Human Services**  
Centers for Disease  
Control and Prevention



# Tdap Vaccine for Preteens and Teens

Last updated JUNE 2014

## Why does my child need Tdap vaccine?

Babies and little kids get shots called DTaP to protect them from diphtheria, tetanus, and pertussis (whooping cough). But as kids get older, the protection from the DTaP shots starts to wear off. This can put your preteen or teen at risk for serious illness. The tetanus-diphtheria-acellular pertussis (Tdap) vaccine is a booster shot that helps protect your preteen or teen from the same diseases that DTaP shots protect little kids from.

- **Tetanus** is caused by a toxin (poison) made by bacteria found in soil. The bacteria enter the body through cuts, scratches, or puncture wounds in the skin. Tetanus can cause spasms which are painful muscle cramps in the jaw muscle (lockjaw) and throughout the body. The spasms can cause breathing problems and paralysis. A preteen or teen with tetanus could spend weeks in the hospital in intensive care. As many as 1 out of 5 people who get tetanus dies.
- **Diphtheria** is not as common as tetanus but can be very dangerous. It spreads from person to person through coughing or sneezing. It causes a thick coating on the back of the nose or throat that can make it hard to breathe or swallow. It can also cause paralysis and heart failure. About 1 out of 10 people who get diphtheria will die from it.
- **Pertussis (whooping cough)** spreads very easily through coughing and sneezing. It can cause a bad cough that makes someone gasp for air after coughing fits. This cough can last for many weeks, which can make preteens and teens miss school and other activities. Whooping cough can be deadly for babies who are too young to have protection from their own vaccines. Often babies get whooping cough from their older brothers or sisters, like preteens or teens, or other people in the family.

## When should my child be vaccinated?

All preteens should get one Tdap shot when they are 11 or 12 years old. If your teen is 13 years old up through 18 years old and hasn't gotten the shot yet, talk to their doctor about getting it for them right away.

## What else should I know about the vaccine?

The Tdap shot has been studied very carefully and is safe. It is recommended by the Centers for Disease Control and Prevention (CDC), the American Academy of Family Physicians, the American Academy of Pediatrics, and the Society for Adolescent Health and Medicine.

The Tdap shot can cause mild side effects, like redness and soreness in the arm where the shot was given, headache, fever, or tiredness. Some preteens and teens might faint after getting the Tdap vaccine or any other shot. To help avoid fainting, preteens and teens should sit or lie down when they get a shot and then for about 15 minutes after getting the shot. Serious side effects from reactions to the Tdap shot are rare.

## How can I get help paying for these vaccines?

The Vaccines for Children (VFC) program provides vaccines for children ages 18 years and younger, who are not insured, Medicaid-eligible, American Indian or Alaska Native. You can find out more about the VFC program by going online to [www.cdc.gov](http://www.cdc.gov) and typing VFC in the search box.


## Where can I learn more?

Your child's doctor or nurse can give you more information about the Tdap vaccine and the other vaccines your child may need. There is also information on CDC's Vaccines for Preteens and Teens website at [www.cdc.gov/vaccines/teens](http://www.cdc.gov/vaccines/teens).

DISTRIBUTED BY:



**U.S. Department of  
Health and Human Services**  
Centers for Disease  
Control and Prevention



## SCPH Clinic

The Immunization Clinic at Summit County Public Health offers immunizations for children, teens and adults. Our skilled Public Health Nurses are ready to help you determine which vaccines are recommended for you or your child. Whether you've just had a child, are preparing your family for the school year, or are simply unsure which vaccines you need, we're ready to help. SCPH offers immunizations at two locations:



### **Summit County Public Health - Fairway Center**

1867 W. Market Street  
Akron, Ohio 44313

**Effective March 5, 2018, the Graham Road Clinic will be open once a week by appointment only. For an appointment, please call 330-375-2772**

### **Summit County Public Health - Graham Road**

1100 Graham Road Circle  
Stow, Ohio 44224

Call today for more information:

[Contact Us \(contact\)](#) 